

Introduction to Graph Theory

From Binary Relations to Feynman Diagrams

Ray D. Sameshima

2025/09/25

About

Graph theory has long served as a powerful tool in physics – from its early role in solving puzzles to its essential use in circuit analysis, modeling state transitions, and representing quantum interactions among fields and particles. This talk presents selected insights from the book [Lectures on Graph Theory – Insights into Feynman Diagrams](#), highlighting how graph-theoretical methods, in combination with linear algebra, form a unified framework for understanding Feynman diagrams and their applications in quantum field theory. We will review the mathematical foundations and classical applications such as Kirchhoff's laws, and then discuss their modern implications in fundamental physics.

Ray D. Sameshima


Lectures on Graph Theory

Insights into Feynman Diagrams

Provides a clear and accessible introduction to mastering Feynman integral computations

Offers a detailed exploration of essential techniques for practical calculations in particle physics

Features numerous examples and algorithms to help readers learn quickly and effectively



© 2025

Brief History

- ▶ Euler – the father of graph theory (1736 Königsberg Bridge Problem)




Figure: The Seven Bridges of Königsberg. (Wikipedia) Can you traverse all seven bridges exactly once?

- ▶ Kirchhoff – the theory of trees (1847)
- ▶ Cayley – chemical problem of trees (1874)
- ▶ Veblen – simplicial complex (1922)
- ▶ Feynman – Feynman diagram (1949)

Graphs are almost everywhere

- ▶ functions
- ▶ function “type” annotations
- ▶ stock-flow diagrams
 - ▶ Equation of Motion (Newton’s second law of motion)
 - ▶ Work-Energy Theorem
 - ▶ Thermodynamics
- ▶ programming languages and data types
- ▶ Markov chain
- ▶ Electric circuits
- ▶ Feynman diagrams

Binary Relations and Functions

Given two sets A, B , a binary relation over A and B is a subset of $A \times B$. If a binary relation $f \subset A \times B$ satisfies additional conditions:

- ▶ for each $a \in A$, there is some $b \in B$ with $(a, b) \in f$,
- ▶ if $(a, b) \in f$ and $(a, b') \in f$, then $b = b'$,

namely, for each $a \in A$ a unique $b \in B$ exists, we write $b = f(a)$ and call f a function from A to B :

$$f: A \rightarrow B.$$

Along with f , $a \in A$ is mapped as $a \xrightarrow{f} b$.


- ▶ We can identify an ordered pair (a, b) of a relation a line from a to b .
- ▶ The type annotation itself is a line from A to B .

Binary Relations and Functions – Example

For a function $f: \mathbb{N} \rightarrow \mathbb{N}$:

$$f(x) = 2x + 1,$$

we can depict:




A function can be identified as a set of rooted trees of height 2.
As functions are everywhere in mathematics, so are graphs.

Graphs and Directed Graphs

A directed graph consists of

- ▶ a set of lines;
- ▶ a set of vertices;
- ▶ an incidence relation.

Pictorially, a graph is the vertices connected with directed lines among them; if the direction is not important, e.g.,



This is a pictorial representation of seven bridges in Königsberg. Euler's negative solution to the problem would be the first discovery of graph theory: **A graph is eulerian – there is a walk that traverses each line exactly once – iff every point has even degree.**

Simpler Examples

- ▶ EoM: The change in linear momentum of a system from \vec{p}_i to \vec{p}_f is the impulse $\vec{f}\Delta t$ on the system: $\vec{p}_i \xrightarrow{\Downarrow \vec{f}\Delta t} \vec{p}_f$.

For a rotational motion: $\vec{L}_i \xrightarrow{\Downarrow \vec{\tau}\Delta t} \vec{L}_f$.

- ▶ WET and TMEC:

The net work on a system is the change in kinetic energy:

$K_i \xrightarrow{\Downarrow W} K_f$. If the work is provided by a “good source,” namely a conservative force, there is an associated potential energy: $P_i \xrightarrow{\Downarrow W} P_f$, and $P_i + K_i = P_f + K_f$.

- ▶ Thermodynamics


- ▶ The work done by an adiabatic process is given by the change in the internal energy: $U_i \xrightarrow{\Downarrow W_{\text{ad.}}} U_f$.

- ▶ The heat flow Q into a thermodynamical system is defined via

$$U_i \xrightarrow[\Downarrow W]{\Downarrow Q} U_f$$


Slightly Involved Examples

A two-state Markov process:




where p represents the transition probability from 1 to 2.

The absolute temperature is defined via the following Carnot cycle:



Electric Circuits

Given the following circuit configuration with EMFs, can we find the currents i_1, i_2, i_3 ?



A day of a graph-theorist

1. Find a mathematical object.
2. Determine the corresponding graph(s).
3. Apply graph-theoretical analyses.

Various Subgraphs

For a given graph, we can extract several important subgraphs such as:

- ▶ Circuits (also known as Loops)
- ▶ (Spanning) Trees
- ▶ Cut-sets

Various Subgraphs

For a given graph, we can extract several important subgraphs such as:

- ▶ Circuits (also known as Loops)
- ▶ (Spanning) Trees
- ▶ Cut-sets

Most graph theorists use personalized terminology in their books, papers, and lectures. ...It is customary to represent a graph by means of a diagram and to refer to it as the graph. (Frank Harary, "Graph Theory")

Circuits – Definition

A circuit of a given graph is a closed and non-self-intersecting walk.

- ▶ Walk

An alternating sequence of vertices and lines

- ▶ Closed walk


A walk with the initial vertex = the final vertex

- ▶ non-self-intersecting


No vertices appear more than once, except for both initial and final vertices

Numbers of Circuits

Can you count the number of circuits, i.e., loops, in an electric circuit:




or that of the corresponding directed graph?




Trees

For simplicity, let us consider connected graphs.

Here is a tree of the underlying graph:



The complement is called the chord-set, or more abstractly, the cotree relative to the tree:



Trees – Definition

A (spanning) tree of a given connected graph is an acyclic \subset -largest subgraph.

Trees – Definition

A (spanning) tree of a given connected graph is an *acyclic* \subset -largest subgraph.

- ▶ *acyclic* : with no circuits

Trees – Definition

A (spanning) tree of a given connected graph is an *acyclic* \subset -*largest* subgraph.

- ▶ *acyclic* : with no circuits
- ▶ *largest* : an additional line makes it cyclic

Trees – Definition


A (spanning) tree of a given connected graph is an *acyclic*

\subset - *largest* subgraph.

► *acyclic* : with no circuits

► *largest* : an additional line makes it cyclic

Removing lines one by one, we can extract a spanning tree.




Trees – Definition

A (spanning) tree of a given connected graph is an *acyclic* \subset - *largest* subgraph.

- ▶ *acyclic* : with no circuits
- ▶ *largest* : an additional line makes it cyclic


Removing lines one by one, we can extract a spanning tree.




*There is one simple and important kind of graph which has been given the same name by all authors, namely a tree.
(Frank Harary, "Graph Theory")*

A fixed tree and associated circuits

For a graph and a spanning tree:




if we recover a deleted line into the spanning tree, we can build a unique loop associated with a spanning tree.




A fixed tree and associated circuits

For a graph and a spanning tree:



if we recover a deleted line into the spanning tree, we can build a unique loop associated with a spanning tree.



A **fundamental circuit** is given by a tree with a line from the cotree.

Cut-sets

A cut-set of a given graph is ...

Cut-sets

A cut-set of a given graph is ... it's complicated!

Duality and Associated Matrices

Algebra is generous: she often gives more than is asked for. (Jean D' Alembert)

Duality and Associated Matrices

Algebra is generous: she often gives more than is asked for. (Jean D' Alembert)

Duality and Associated Matrices

Linear Algebra is generous: she often gives more than is asked for.

Duality and Associated Matrices

Linear Algebra is generous: she often gives more than is asked for. (Ray D. Sameshima)

Associated Matrix Representations are better for algebraic and computer processing than pictorial representations:

- ▶ Incidence matrix \mathcal{A}

Vertices vs. Lines

For a vertex a and a line l , $\mathcal{A}_{a,l}$ is $+1$ iff l is from a , -1 iff l is to a , 0 otherwise.

- ▶ Circuit matrix \mathcal{C}

Circuits vs. Lines

- ▶ Cut-set matrix \mathcal{S}

Cut-sets vs. Lines

Linear Algebra is generous

Let G be a connected graph with M vertices and N lines. Then the incidence matrix is $M \times N$. We call $r = M - 1$ the rank of the graph G , and $\mathbb{b}_1 = N - r$ the nullity. With a fixed vertex a and a fixed spanning tree T in G , if we write

$$\mathcal{A}^a = (\mathcal{A}_{T^*}^a \quad \mathcal{A}_T^a) \quad (4.79)$$

then the circuit matrix \mathcal{C}_{T^*} and the cut-set matrix \mathcal{S}_T can be written as

$$\mathcal{C}_{T^*} = (1_{\mathbb{b}_1} \quad \mathcal{B}_T), \quad \mathcal{S}_T = (-\mathcal{B}_T^\top \quad 1_r) \quad (4.75)$$

and

$$\mathcal{B}_T^\top = -(\mathcal{A}_T^a)^{-1} \mathcal{A}_{T^*}^a. \quad (4.81)$$

Linear Algebra is generous

Let G be a connected graph with M vertices and N lines. Then the incidence matrix is $M \times N$. We call $r = M - 1$ the rank of the graph G , and $\mathfrak{b}_1 = N - r$ the nullity. With a fixed vertex a and a fixed spanning tree T in G , if we write

$$\mathcal{A}^a = (\mathcal{A}_{T^*}^a \quad \mathcal{A}_T^a) \quad (4.79)$$

then the circuit matrix \mathcal{C}_{T^*} and the cut-set matrix \mathcal{S}_T can be written as

$$\mathcal{C}_{T^*} = (1_{\mathfrak{b}_1} \quad \mathcal{B}_T), \quad \mathcal{S}_T = (-\mathcal{B}_T^\top \quad 1_r) \quad (4.75)$$

and

$$\mathcal{B}_T^\top = -(\mathcal{A}_T^a)^{-1} \mathcal{A}_{T^*}^a. \quad (4.81)$$

One of the bridging tools is the *rank-nullity* theorem.

Electric Circuit Analysis

Let $G = l_1 \left(\begin{array}{c} \text{a} \\ \uparrow l_2 \\ \text{b} \end{array} \right) l_3$ be an underlying directed graph. If

we select a spanning tree $T = \begin{array}{c} \text{a} \\ \text{b} \end{array} l_3$ and the corresponding

cotree $T^* = l_1 \left(\begin{array}{c} \text{a} \\ \uparrow l_2 \\ \text{b} \end{array} \right)$, we have the following “fundamental”

circuits and a cut-set:

$$C_1 = l_1 \left(\bigcirc \right) l_3, C_2 = l_2 \uparrow \left(\bigcap \right) l_3, S_1 = \left(\begin{array}{c|c} l_1 & l_2 \end{array} \right) l_3$$

Electric Circuit Analysis (continued)

I.e., this graph and the electric circuit have two “loops.” If we assign the loop currents j_1 and j_2 along the loops l_1 and l_2 , respectively, Kirchhoff's junction rule is automatically guaranteed. Then the corresponding line currents are

$$i_1 = j_1, \quad i_2 = j_2, \quad i_3 = j_1 + j_2.$$

Kirchhoff's loop rule becomes the following 2×2 system:

$$0 = R_1 j_1 + (-E_2 + R_3 (j_1 + j_2))$$

$$0 = (R_2 j_2 - E_1) + (-E_2 + R_3 (j_1 + j_2))$$

That is, the linear dimension of the electrical problem is **the nullity** of the graph:

$$\mathfrak{b}_1 = N - M + 1.$$

Electric Circuit Analysis (continued)

I.e., this graph and the electric circuit have two “loops.” If we assign the loop currents j_1 and j_2 along the loops l_1 and l_2 , respectively, Kirchhoff’s junction rule is automatically guaranteed. Then the corresponding line currents are

$$i_1 = j_1, \quad i_2 = j_2, \quad i_3 = j_1 + j_2.$$

Kirchhoff’s loop rule becomes the following 2×2 system:

$$0 = R_1 j_1 + (-E_2 + R_3 (j_1 + j_2))$$

$$0 = (R_2 j_2 - E_1) + (-E_2 + R_3 (j_1 + j_2))$$

That is, the linear dimension of the electrical problem is **the nullity** of the graph:

$$\mathbb{b}_1 = N - M + 1.$$

Moreover, it is the number of **fundamental circuits** in the graph.

Electric Circuit Analysis (continued)

Furthermore, one can show the following power conservation – instantaneous energy conservation – known as Tellegen's theorem:

$$\mathcal{I}^\top \mathcal{V} = 0, \quad (4.99)$$


where \mathcal{I} is the current vector across the electrical circuit and \mathcal{V} is the potential difference:

Among the theorems of network theory Tellegen's theorem is unusual in that it depends solely upon Kirchhoff's laws and the topology of the network. (Penfield, P. Spence, R. and Duinker, S. "Tellegen's Theorem and Electrical Networks")

Feynman Diagrams?

A graphical representation of the quantum interactions among particles and fields.

A Feynman diagram consists of a graph with external legs:



The external legs, the dashed lines, represent interactions with the outside world, like ports in an electric circuit.

Feynman Diagrams 101

They are graded with respect to the number of circuits in the graph parts; the number of circuits is the order of the asymptotic expansion of S -matrix:

zero-loop diagrams + one-loop diagrams + two-loop diagrams + \dots

Each term in the expansion is an involved integral, represented as a Feynman diagram. To achieve higher precision, we require more diagrams, which makes the computation challenging in both memory space and processing speed.

Using the graph-theoretical tools, we have a systematic way to write the integral with polynomials associated with the underlying Feynman diagram.

Not only the numerical values but also their analytical properties of integrals are of interest.

Feynman Diagrams 101

They are graded with respect to the number of circuits in the graph parts; the number of circuits is the order of the asymptotic expansion of S -matrix:

zero-loop diagrams + one-loop diagrams + two-loop diagrams + \dots
tree-level

Each term in the expansion is an involved integral, represented as a Feynman diagram. To achieve higher precision, we require more diagrams, which makes the computation challenging in both memory space and processing speed.

Using the graph-theoretical tools, we have a systematic way to write the integral with polynomials associated with the underlying Feynman diagram.

Not only the numerical values but also their analytical properties of integrals are of interest.

Feynman Diagrams 101

They are graded with respect to the number of circuits in the graph parts; the number of circuits is the order of the asymptotic expansion of S -matrix:

zero-loop diagrams + one-loop diagrams + two-loop diagrams + \dots
tree-level quantum corrections

Each term in the expansion is an involved integral, represented as a Feynman diagram. To achieve higher precision, we require more diagrams, which makes the computation challenging in both memory space and processing speed.

Using the graph-theoretical tools, we have a systematic way to write the integral with polynomials associated with the underlying Feynman diagram.

Not only the numerical values but also their analytical properties of integrals are of interest.

Feynman Diagrams 101

They are graded with respect to the number of circuits in the graph parts; the number of circuits is the order of the asymptotic expansion of S -matrix:

zero-loop diagrams + one-loop diagrams + two-loop diagrams + \dots
tree-level quantum corrections

Each term in the expansion is an involved integral, represented as a Feynman diagram. To achieve higher precision, we require more diagrams, which makes the computation challenging in both memory space and processing speed.

Using the graph-theoretical tools, we have a systematic way to write the integral with **polynomials associated with the underlying Feynman diagram**.

Not only the numerical values but also their analytical properties of integrals are of interest.

An Open Problem

Similar to electric currents across lines in an electric circuit, each line in a Feynman diagram has the corresponding energy-momentum flow.

Question: Is there any voltage-like quantity associated with a Feynman diagram?

If such a quantity exists, thanks to Tellegen's theorem, we may obtain yet another quantum conservation law!

Part of the book series:
Lecture Notes in Physics



Get 20% off with this code: **SPRAUT**

Available on Springer Nature Link



[link.springer.com/book/
9783031822186](https://link.springer.com/book/9783031822186)

Please note that promotional coupons are only valid for English-language Springer, Apress, and Palgrave Macmillan books & eBooks and are redeemable on link.springer.com only. Titles affected by fixed book price laws, forthcoming titles and titles temporarily not available on Springer Nature Link are excluded from promotions, as are reference works, handbooks, encyclopedias, subscriptions, or bulk purchases. The currency in which your order will be invoiced depends on the billing address associated with the payment method used, not necessarily your home currency. Regional VAT/tax may apply. Promotional prices may change due to exchange rates. Promotions are valid for individual customers only. Booksellers, book distributors, and institutions such as libraries and corporations please visit springernature.com/contact-us. Promotions do not work in combination with other discounts or gift cards.

link.springer.com/book/9783031822186